

JAMA Otolaryngology-Head & Neck Surgery | Original Investigation

Addition of Hyperbaric Oxygen Therapy vs Medical Therapy Alone for Idiopathic Sudden Sensorineural Hearing Loss

A Systematic Review and Meta-analysis

Tae-Min Rhee, MD, DMO/UMO; Doyeon Hwang, MD; Jee-Soo Lee, MD; Jonghanne Park, MD, PhD; Joo Myung Lee, MD, MPH, PhD

IMPORTANCE Sudden sensorineural hearing loss (SSNHL) causes substantial disease burden for both individuals and socioeconomic aspects. The benefit of hyperbaric oxygen therapy (HBOT) in addition to standard medical therapy (MT) for idiopathic SSNHL has been unclear.

OBJECTIVE To perform a systematic review and meta-analysis to compare HBOT + MT with MT alone as a treatment for patients with SSNHL.

DATA SOURCES PubMed, Embase, and the Cochrane Database of Systematic Reviews were systematically searched up to February 2018.

STUDY SELECTION Randomized clinical trials and nonrandomized studies comparing HBOT + MT with MT alone for SSNHL treatment.

DATA EXTRACTION AND SYNTHESIS Two investigators independently screened the eligible studies, established data, and assessed quality and risk of bias. A systematic review and meta-analysis using random-effects models was conducted.

MAIN OUTCOMES AND MEASURES The primary outcome was complete hearing recovery, and secondary outcomes were any hearing recovery and absolute hearing gain.

RESULTS Three randomized clinical trials and 16 nonrandomized studies comparing outcomes after HBOT + MT vs MT alone in 2401 patients with SSNHL (mean age, 45.4 years; 55.3% female) were included. Pooled odds ratios (ORs) for complete hearing recovery and any hearing recovery were significantly higher in the HBOT + MT group than in the MT alone group (complete hearing recovery OR, 1.61; 95% CI, 1.05-2.44 and any hearing recovery OR, 1.43; 95% CI, 1.20-1.67). Absolute hearing gain was also significantly greater in the HBOT + MT group than in the MT alone group. The benefit of HBOT was greater in groups with severe to profound hearing loss at baseline, HBOT as a salvage treatment, and a total HBOT duration of at least 1200 minutes.

CONCLUSIONS AND RELEVANCE The addition of HBOT to standard MT is a reasonable treatment option for SSNHL, particularly for those patients with severe to profound hearing loss at baseline and those who undergo HBOT as a salvage treatment with a prolonged duration. Optimal criteria for patient selection and a standardized regimen for HBOT should be applied in routine practice, with future trials to investigate maximal treatment benefit.

JAMA Otolaryngol Head Neck Surg. 2018;144(12):1153-1161. doi:10.1001/jamaoto.2018.2133
Published online September 27, 2018.

[+ Supplemental content](#)

[+ CME Quiz at \[jamanetwork.com/learning\]\(http://jamanetwork.com/learning\) and CME Questions page 1192](#)

Author Affiliations: Department of Undersea and Hyperbaric Medicine, Maritime Medical Research Center, National Maritime Medical Center, Changwon, Republic of Korea (Rhee); Department of Internal Medicine, Seoul National University Hospital, Seoul, Republic of Korea (Rhee, Hwang, Park); Department of Laboratory Medicine, Hallym University Sacred Heart Hospital, Anyang, Republic of Korea (J.-S. Lee); Department of Internal Medicine and Cardiovascular Center, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Republic of Korea (J. M. Lee).

Corresponding Author: Tae-Min Rhee, MD, DMO/UMO, Department of Undersea and Hyperbaric Medicine, Maritime Medical Research Center, National Maritime Medical Center, 963-gil 35, Jinhaedae-ro, Jinhae-gu, Changwon, Gyeongsangnam-do 51640, Republic of Korea (imcrtm@gmail.com).

Sudden sensorineural hearing loss (SSNHL) is a subset of sudden hearing loss that occurs within 72 hours and is defined as a hearing loss of at least 30 dB identified at 3 or more consecutive frequencies.¹ Such hearing loss is not uncommon, with an incidence of 5 to 20 cases per 100 000 in the general population.¹ It causes severe discomfort because of sudden deafness on 1 or both sides and increases the risk of accidents caused by decreased spatial perception.² This risk further results in substantial disease burden not only among individuals but also socioeconomically.³ However, SSNHL is a disease entity from which recovery by prompt and appropriate treatment could dramatically improve patient quality of life and reduce the need for hearing aids.³

Corticosteroids, antiviral agents, vasodilators, and hyperbaric oxygen therapy (HBOT) are the currently available treatment options for SSNHL, but their comparative efficacy is unclear.¹ To date, the most widely used treatment for SSNHL is systemic and/or intratympanic corticosteroids.¹ As another option, HBOT is a treatment that may relieve edema and ischemia by administering high-pressure oxygen into the inner ear to restore hearing.⁴ Since the first case report⁵ of HBOT for SSNHL treatment in the 1960s, a number of randomized clinical trials (RCTs)⁶⁻⁸ and nonrandomized studies⁹⁻²⁴ investigating the benefit of HBOT have been reported. As an adjunctive treatment to standard medical therapy (MT) that includes systemic corticosteroids, HBOT has been found to promote hearing gain.^{25,26} However, there is limited evidence that HBOT definitively improves the outcome of SSNHL.²⁵ Therefore, we performed a systematic review and meta-analysis of 19 studies to compare HBOT + MT with MT alone as a treatment for patients with SSNHL.

Methods

Detailed study methods are available in the eMethods in the [Supplement](#); the search strategy in PubMed, Embase, and Cochrane Database of Systematic Reviews is described, as well as the characteristics of the excluded studies.

Data Sources and Searches

PubMed, Embase, and the Cochrane Database of Systematic Reviews were systematically searched up to February 2018 for published or unpublished studies. This electronic search strategy was augmented by a manual examination of references cited in articles, recent reviews, editorials, and meta-analyses. No restrictions were imposed on the language, study period, or sample size. The search strategy is described in detail in the eMethods in the [Supplement](#).

Study Selection and Outcome Definition

Studies meeting each of the following criteria were eligible for the meta-analysis: (1) performed before February 2018; (2) definitively reported outcomes, including complete hearing recovery and any hearing recovery or absolute hearing gain; (3) compared outcomes after HBOT + MT vs MT alone; (4) included systemic and/or intratympanic corticosteroids in the MT protocol; and (5) used a clear definition of SSNHL.

Key Points

Question What is the benefit of the addition of hyperbaric oxygen therapy (HBOT) vs medical therapy alone for sudden sensorineural hearing loss (SSNHL)?

Findings Results of this meta-analysis including 2401 patients with SSNHL significantly favored HBOT plus standard medical therapy (MT) over MT alone for complete hearing recovery and any hearing recovery, as well as for absolute hearing gain. The benefit of HBOT was greater in groups with severe hearing loss at baseline, HBOT as a salvage treatment, and a total HBOT duration of at least 1200 minutes.

Meaning The benefit of HBOT for SSNHL may be greater for those who had severe hearing loss at baseline or who failed to recover after MT; optimal criteria for patient selection and a standardized regimen for HBOT should be established in future trials.

Studies reporting SSNHL treatment outcomes without comparator or control groups were excluded. Two investigators (T.-M.R. and D.H.) independently screened titles and abstracts, identified duplicates, reviewed full articles, and determined their eligibility. Disagreements were resolved by discussion. The last search was performed in February 2018. The primary outcome was complete hearing recovery, and secondary outcomes were any hearing recovery and absolute hearing gain.

Data Extraction and Quality Assessment

Data were compiled for the meta-analysis using a standardized form to extract the following characteristics of studies: study design, number of patients, treatment protocol (HBOT + MT or MT alone), outcome definitions, treatment protocols, and patient demographics. The quality of the eligible studies was assessed using the Cochrane Collaboration's tool for assessing risk of bias for RCTs. The Newcastle-Ottawa Scale (NOS) and the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist were used to assess the quality of nonrandomized prospective studies. However, the thresholds of the NOS or STROBE checklist scores were not grounds for individual study exclusion.

Data Synthesis and Statistical Analysis

Random-effects models were applied for primary and secondary outcome analyses, and odds ratios (ORs) are presented with 95% CIs in the statistical summaries. Because the selected studies were heterogeneous in terms of study population and protocol, fixed-effects models were used for sensitivity analyses, and the results yielded were checked for similarity. The pooled ORs and 95% CIs of the random-effects and fixed-effects models were calculated using the restricted maximum likelihood and Mantel-Haenszel methods, respectively.²⁷ The pooled results for the continuous variables are presented as the weighted mean difference with a 95% CI between the HBOT + MT group and the MT alone group.

Statistical heterogeneity was quantified using I^2 statistics. Publication bias, a known threat to meta-analysis validity created by the preferential publishing of studies with statistically significant or clinically favorable results,²⁸ was

assessed through funnel plot asymmetry using Egger test and Begg test. If visible asymmetry of funnel plots was observed, the trim-and-fill method was used to estimate the number of missing studies and calculate the corrected risk ratio as if such studies were present. Exploratory meta-regressions were performed to assess the association between effect size (log ORs) and the mean age, proportion of men, and initial hearing level.

Hearing recovery rates were separately analyzed according to the severity of the initial hearing loss. Subgroup analyses were used to assess differential associations with the following methods: (1) the statistical model (fixed effects vs random effects), (2) the HBOT strategy (salvage treatment vs adjunctive treatment), (3) the total HBOT duration (≥ 1200 vs < 1200 minutes), (4) the maximal pressure during HBOT (≥ 2.5 vs < 2.5 atmospheric absolute pressure [ATA]), and (5) the response assessment point (≥ 3 vs < 3 months after treatment). Two-sided $P < .05$ was considered statistically significant. Statistical computations were performed with a standard software program (Stata/SE, version 12.0; Stata-Corp LP). The present study complied with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (eTable 1 in the Supplement) and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines.

Results

Search Findings

A total of 194 citations were identified. Among these, 24 articles were retrieved for a full review, and 19 met the inclusion criteria (Figure 1). The characteristics of the 5 excluded studies after a full article review are detailed in the eMethods in the Supplement. The final 19 studies included 2401 patients with SSNHL (mean age, 45.4 years; 55.3% female) grouped by treatment protocol as HBOT + MT (1055 of 2401 [43.9%]) or MT alone (1346 of 2401 [56.1%]). Among the 19 target studies, 14 provided the rates of complete hearing recovery and any hearing recovery, and 11 provided absolute hearing gain.


Study Characteristics and Risk of Bias

Within Studies

The main characteristics of the individual studies are summarized in the Table. Most studies were retrospective observational investigations, while 2 were prospective studies and 3 were RCTs. All studies exclusively enrolled patients with SSNHL who were undergoing treatment. Follow-up duration varied from weeks to months, with 8 studies yielding outcomes immediately after treatment. The mean onset to HBOT time ranged from 3 to 48.2 days, and HBOT was used as a salvage treatment in 7 studies. Demographic features, including age and sex, were evenly distributed in all studies. The initial hearing level ranged from 43.6 to 86.8 dB.

eTable 2 and eTable 3 in the Supplement summarize the assessed risk of bias by study design. Two RCTs did not report any random sequence generation method. Although no masking attempt was observed, all studies objectively

Figure 1. Flow Diagram of Study Selection



The flow diagram is shown according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

defined clinical end points, with a complete follow-up. Therefore, the lack of masking had a low probability of influencing outcomes. All nonrandomized studies met at least 17 variables of the STROBE checklist and fulfilled the adequacy criteria of the NOS for nonrandomized studies.

Addition of HBOT and Hearing Recovery and Absolute Hearing Gain

The rates of complete hearing recovery in the HBOT + MT and MT alone groups were 264 of 897 (29.4%) and 241 of 1167 (20.7%), respectively, and the rates of any hearing recovery were 621 of 919 (67.6%) and 585 of 1194 (49.0%), respectively. The pooled results from random-effects models significantly favored the HBOT + MT group over the MT alone group for complete hearing recovery (pooled OR, 1.61; 95% CI, 1.05-2.44) and for any hearing recovery (pooled OR, 1.43; 95% CI, 1.20-1.67) (Figure 2). Significant heterogeneity was observed for both outcomes. Funnel plots, along with Egger test and Begg test results, demonstrated no significant publication bias for complete hearing recovery and any hearing recovery, and there was no need for trim-and-fill adjustment because of the absence of asymmetry (eFigure 1 in the Supplement).

The weighted mean differences of absolute hearing gain are presented as the mean value for the overall frequencies, as well as for each frequency level (eFigure 2 in the Supplement). Absolute hearing gain was significantly greater in the HBOT + MT group than in the MT alone group for the overall frequencies


Table. Characteristics of Studies Selected for Meta-analysis^a

Source	Study Period	Study Design	No. of Patients		Complete Hearing Recovery >50%	Any Hearing Recovery >25%	Protocol	HBOT	Follow-up Duration	Onset to HBOT Time	HBOT as a Salvage Treatment
			HBOT + MT	MT Alone							
Cavallazzi et al, ²² 1996	NR	Retrospective	34	30	Recovery >50%	Recovery >25%	MT Betamethasone, dose NR	2.5 ATA for 60 min, 15 sessions	After treatment	NR	No
Aslan et al, ²³ 2002	1995-1999	Retrospective	25	25	NA	NA	Prednisone at an initial dose of 1 mg/kg/d for 2 wk	2.4 ATA for 100 min, 20 sessions	8 wk	5.8 d	No
Narozny et al, ⁹ 2006	1980-2000	Retrospective	52	81	NA	NA	Prednisone 30 mg by mouth per day in decreasing dose for up to 14 d	2.5 ATA for 60 min, 16 ± 6 sessions	After treatment	NR	No
Topuz et al, ⁶ 2004	1998-2002	RCT	34	21	NA	NA	Prednisone at an initial dose of 1 mg/kg per day for 2 wk	2.5 ATA for 90 min, 25 sessions	4 wk	<2.0 wk	No
Desloovere et al, ¹⁰ 2006	NR	Retrospective	56	160	AHG>20 dB	AHG>10 dB	Tapered dose of hydrocortisone starting at 250 mg IV	1.5 ATA for 60 min, 14 sessions	13.4 mo	48.2 d	Yes
Satar et al, ¹¹ 2006	1996-2003	Retrospective	37	17	MHL<25 dB	AHG>10 dB	Dexamethasone 4 mg IV twice a day for 7 d	2.5 ATA for 90 min, 18 sessions	After treatment	5.0 d	No
Dundar et al, ¹² 2007	2002-2004	Prospective	55	25	MHL<25 dB	AHG>15 dB	Dexamethasone 4 mg IV twice a day for 7 d	2.8 ATA for 90 min, 10-28 sessions	After treatment	<1.0 wk	No
Fujimura et al, ¹³ 2007	1979-2005	Retrospective	67	63	MHL<20 dB	AHG>10 dB	Dexamethasone 8 mg IV, followed by tapered doses for 12 d	2.5 ATA for 60 min, 10 sessions	1 mo	6.4 d	No
Cekin et al, ⁷ 2009	1994-2006	RCT	36	21	AHG>50 dB	AHG>10 dB	Prednisolone 1 mg/kg starting dose, tapering in 3 wk	2.5 ATA for 90 min, 10 sessions	After treatment	3.0 d	No
Ohno et al, ¹⁴ 2010	2001-2008	Retrospective	48	44	AHG>30 dB	AHG>10 dB	Betamethasone 10 mg/d tapered for 10 d or prednisolone 30 mg/d tapered for 12 d	2.0 ATA for 60 min, 10 sessions	23 wk	7.4 wk	Yes
Alimoglu et al, ¹⁵ 2011	2004-2010	Retrospective	61	58	MHL<25 dB	AHG>15 dB	Prednisolone 1 mg/kg or equivalent, 10-mg taper every 3 d	2.5 ATA for 120 min, 20 sessions	After treatment	<1.0 mo	No
Liu et al, ¹⁶ 2011	1999-2009	Retrospective	112	353	<15 dB within unaffected ear	AHG>10 dB	Betamethasone 12 mg tapered for 6 d, followed by oral prednisolone 20 mg/d for 8 d	2.5 ATA for 60 min, 10-20 sessions	180 d	<2.0 wk	No
Cvorovic et al, ⁸ 2013	2005-2011	RCT	25	25	NA	NA	Dexamethasone 40 mg IV for 3 d, followed by 10 mg/d for 3 d	2.0 ATA for 60 min, 20 sessions	After treatment	<4.0 wk	Yes
Yang et al, ¹⁷ 2013	NR	Retrospective	22	27	AHG>15 dB	AHG>10 dB	Dexamethasone 20 mg IV for 2 d, tapered for 5 d, followed by oral prednisolone 30 mg/d for 5 d	2.5 ATA for 120 min, 10 sessions	2 mo	4.2 d	Yes
Capuano et al, ¹⁸ 2015	2010-2013	Retrospective	100	100	<15 dB within unaffected ear	AHG>10 dB	Methylprednisolone 40 mg IV for 7 d, 20 mg for another 3 d	2.5 ATA for 90 min, 16 sessions	180 d	<90.0 d	No
Pezzoli et al, ¹⁹ 2015	2011-2013	Prospective	23	21	<15 dB within unaffected ear	AHG>10 dB	Betamethasone 4 mg/d for 6 d, salvage with dexamethasone 25 mg by mouth for 7 d	2.5 ATA for 30 min, 15 sessions	3 mo	9.9 d	Yes
Psillas et al, ²⁰ 2015	2013-2015	Retrospective	15	30	<15 dB within unaffected ear	AHG>10 dB	Dexamethasone 24 mg/d IV, tapered for 1 wk	2.2 ATA for 90 min, 15 sessions	3 mo	24.0 d	Yes
Hosokawa et al, ²¹ 2017	2011-2015	Retrospective	167	160	MHL<20 dB	AHG>10 dB	Prednisolone 80 mg or hydrocortisone 400 mg tapered over 10 d	1.5 ATA for 60 min, 10 sessions	3 mo	<1.0 mo	Yes
Ricciardiello et al, ²⁴ 2017	2009-2016	Retrospective	86	85	MHL<25 dB	AHG>15 dB	Dexamethasone 8 mg/d or methylprednisolone 1 mg/kg tapered for 15 d	2.5 ATA for 90 min, 15-21 sessions	After treatment	7.0 d	No

^a Demographics of the overall population are available in eTable 4 in the Supplement.

Abbreviations: AHG, absolute hearing gain; ATA, atmospheric absolute pressure; HBOT, hyperbaric oxygen therapy; IV, intravenously; MHL, mean hearing level; MT, medical therapy; NA, not applicable; RCT, randomized clinical trial.

Figure 2. Benefit of the Addition of Hyperbaric Oxygen Therapy (HBOT) on Hearing Recovery



(weighted mean difference, 8.74; 95% CI, 5.05-12.43 dB). Similar trends were observed for each frequency level, although none were statistically significant except at 500 Hz.

Severity of the Initial Hearing Loss and Hearing Recovery

To assess the differential association of HBOT according to the severity of the initial hearing loss, the outcomes were compared separately by groups with severe to profound hearing loss (≥ 70 dB) or mild to moderate hearing loss (< 70 dB) (Figure 3). For complete hearing recovery and any hearing recovery, HBOT + MT was more beneficial than MT alone in the group with severe to profound hearing loss than in the group with mild to moderate hearing loss.

Subgroup Analyses

In a subgroup analysis for complete hearing recovery, the benefit of the strategy and protocols of HBOT on the outcome was apparent (Figure 4). The OR for complete hearing recovery significantly favored HBOT + MT over MT alone, especially for those who underwent HBOT as a salvage treatment and had a total HBOT duration of at least 1200 minutes. The higher maximal pressure during HBOT of at least 2.5 ATA was not beneficial for complete hearing recovery. Among other subgroups of the statistical model and the time point of response assessment, the benefit of HBOT on complete hearing recovery was similar. For any hearing recovery, the favorable results of the HBOT + MT group compared

with the MT alone group were consistent across all subgroups (eFigure 3 in the Supplement).

Demographic Characteristics and Baseline Hearing Level and Hearing Recovery

To evaluate the association of age and sex with hearing recovery, exploratory meta-regression analyses were conducted. The mean patient age in each study was not significantly associated with a benefit of HBOT on hearing recovery (eFigure 4 in the Supplement). A significant association between sex and the OR for any hearing recovery was observed, demonstrating that a lower proportion of men was associated with a greater benefit of HBOT (eFigure 5 in the Supplement).

Discussion

We performed a meta-analysis comprising 2401 patients with SSNHL in 19 studies that compared the hearing recovery rate and absolute hearing gain between HBOT + MT and MT alone groups. The principal findings were as follows: (1) The pooled ORs for

complete hearing recovery and any hearing recovery were significantly higher in the HBOT + MT group. (2) Absolute hearing gain was also significantly greater in the HBOT + MT group than in the MT alone group, showing similar trends at all individual frequencies. (3) The HBOT appeared to be more beneficial in patients with severe to profound hearing loss at baseline. (4) The results were consistent across various subgroups, and the benefit of HBOT was prominent in groups with HBOT as a salvage treatment and a total HBOT duration of at least 1200 minutes.

Prior Evidence on HBOT for SSNHL

The treatment of SSNHL to date has focused mainly on the improvement of blood flow and increased oxygen supply in the inner ear. Although there are several treatment options, including systemic and/or intratympanic corticosteroids, antiviral agents, vasodilators, anticoagulants, and plasma expanders, no definitive treatment for SSNHL has been proved through qualified trials or meta-analysis.²⁵ Because corticosteroids are thought to reduce the inflammation and edema associated with SSNHL,²⁹ experts recommend systemic and/or intratympanic corticosteroids first unless specific contraindications are present.³

Figure 3. Hearing Recovery Rate According to the Severity of the Initial Hearing Loss





Figure 4. Subgroup Analyses for Complete Hearing Recovery



Benefit of hyperbaric oxygen therapy (HBOT) on complete hearing recovery according to the various subgroups is shown. ATA indicates atmospheric absolute pressure; MT, medical therapy; OR, odds ratio.

The efficacy of HBOT was updated in the Cochrane Database of Systematic Reviews in 2012 by Bennett et al.²⁵ In their review, a 25% chance of hearing recovery was significantly higher in the HBOT group (risk ratio, 1.39; 95% CI, 1.05-1.84), while there was no difference in a 50% chance of hearing recovery (risk ratio, 1.53; 95% CI, 0.85-2.78). A greater improvement of the mean threshold in the HBOT group than in the control group was also observed (mean difference, 15.6; 95% CI, 1.5-29.8 dB). The use of systemic and/or intratympanic corticosteroids plus HBOT has been considered the most effective SSNHL treatment for several reasons.³ In the study by Bennett et al,²⁵ only 392 patients in 7 studies were analyzed; because the outcome was divided into absolute gains and hearing recovery rates, the statistical power in their review was insufficient, and the results are likely unreliable. In addition, the included studies provided no detailed treatment strategy or protocol information, making it impossible to perform subgroup analyses according to the severity of the initial hearing loss or HBOT protocols.

Since that review, several studies have been published presenting controversial results regarding the benefit of HBOT. Saesen et al²⁶ reported in a recent narrative review that HBOT has a positive benefit on hearing gain as an adjunctive therapy to standard MT. Their study also indicated that HBOT could have a positive role as a salvage treatment after the failure of initial corticosteroid treatment. Therefore, a comprehensive meta-analysis covering recent studies was necessary.

HBOT in Patients With SSNHL

In our meta-analysis, HBOT was found to provide a significant benefit as an additional treatment option, along with systemic and/or intratympanic corticosteroids. In particular, this study demonstrates for the first time to date that HBOT + MT is associated with a significant improvement in complete hearing recovery and in any hearing recovery compared with MT alone. This finding is important evidence for the clinical implications of HBOT in SSNHL treatment.

To date, hypotheses about the benefit of HBOT have been suggested based on various theoretical backgrounds. The structures in the cochlea are vulnerable to a decrease in tissue oxygen supply.³ In addition, the supply to the cochlea depends on oxygen diffusion through the capillaries rather than direct vascular oxygenation.³⁰ Therefore, it is thought that the reduction of blood flow to the inner ear and resultant ischemia are the most important mechanisms for the occurrence of idiopathic SSNHL.³¹ Using HBOT, it is possible to maximize the oxygen partial pressure supplied to the inner ear.²⁵ This process can minimize ischemic damage after SSNHL and aid vascular recovery.³² Furthermore, it can provide antibacterial effects through oxygen radicals and promote angiogenesis with tissue regeneration.⁴ The present study supports these theories by incorporating updated clinical data, providing valuable evidence that HBOT + MT is the most beneficial treatment option for SSNHL.

In terms of safety, serious complications from HBOT are uncommon.²⁵ Although middle ear, sinus, and pulmonary barotrauma can occur, their incidence is low. Only ear fullness, which is completely recoverable, is known to be a com-

mon adverse effect.³³ The use of HBOT is limited in that it is only available in special facilities, and patients with claustrophobia cannot undergo HBOT.³² However, patients with a history of pneumothorax or those with claustrophobia can be excluded from HBOT treatment at the initial patient interview. Therefore, the benefit of hearing gain through HBOT may exceed any possible harm, particularly if there has been no response to standard treatment with corticosteroids or if there is severe initial hearing loss.

In South Korea, HBOT treatment costs US \$100 per session, but the actual patient contribution is US \$40 based on nationwide medical insurance coverage. In the United States, it costs approximately US \$300 per session. However, considering the adverse effects of idiopathic SSNHL on patient quality of life and the risk of accidents caused by decreased spatial perception, HBOT as a salvage treatment may be considered at least for refractory cases after medical treatment with corticosteroids. To achieve better cost-effectiveness, well-defined indications for HBOT and standardized HBOT regimens should be established and applied.

Clinical Implications From Subgroup Analyses

The results of earlier studies^{6,16} have indicated that the benefit of HBOT may be greater in more severely affected patients. Experts have also suggested that HBOT may be more effective at ages younger than 60 years, with early HBOT application within 3 months, and in a group with greater initial hearing loss of more than 60 dB, although the level of evidence is weak.¹ However, the Cochrane Database of Systematic Reviews in 2012 concluded that the influence of the severity of the initial hearing loss could not be confirmed.²⁵ In the present meta-analysis, we found that the benefit of the addition of HBOT was greater in the group with severe to profound hearing loss of at least 70 dB. However, because there were limited studies that provided separate outcomes based on initial severity, the statistical power was low, and follow-up studies are required.

Furthermore, the group with HBOT as a salvage treatment received greater benefit than the group with HBOT as an adjunctive treatment, which is also controversial and warrants further trials that control for the HBOT strategy. Our meta-regression results did not show any evidence that age influences the benefit of HBOT. Furthermore, because HBOT was performed within 3 months in all enrolled participants, the differential association according to the onset to HBOT time could not be assessed.

It is generally recommended that 100% oxygen at 2.0 to 2.5 ATA should be administered for 10 to 20 days, with a 90-minute session each day,³ but there are no universal HBOT protocols that have been proved to be effective. For the first time to date, we have analyzed the associations of the duration and maximal pressure of HBOT with a treatment benefit. Particularly for complete hearing recovery, HBOT + MT was favored in the group with a total HBOT duration of at least 1200 minutes. However, the maximal pressure during HBOT was not found to have a significant association with any benefit of HBOT. These results are expected to have an important role in the standardization of clinical indications for and proto-

cols of HBOT. Sudden sensorineural hearing loss with severe to profound hearing loss (≥ 70 dB) at baseline can be considered an appropriate indication for applying HBOT. Given the greater benefit when HBOT is used as a salvage treatment, HBOT may be applied to refractory cases that do not respond to medical treatment 2 to 4 weeks after the onset of hearing loss. In terms of HBOT strategy, a 2-week protocol with a 90-minute session per day may be recommended or alternatively a 20-day protocol with a 60-minute session per day to achieve a total HBOT duration of at least 1200 minutes. However, because increasing maximal pressure during HBOT to 2.5 or higher ATA did not provide any benefit for hearing recovery, the air pressure may be kept at 2.0 to 2.5 ATA as currently recommended.

Future trials should use a standardized HBOT regimen to investigate the efficacy of HBOT as a salvage treatment for patients refractory to MT with corticosteroids. In particular, the appropriate intervals should be clarified between onset, assessment of initial treatment failure, and HBOT as a salvage treatment.

Limitations

This meta-analysis has several limitations. First, the selected studies varied in their clinical and methodological characteristics. Second, although there was no evidence of publication bias, the main results showed considerable heterogeneity. Third, considering that most studies except RCTs did not pro-

vide adjusted results, measured or unmeasured confounder effects could exist because of different baseline characteristics between study and control groups. Fourth, because a substantial proportion of patients with SSNHL experience spontaneous recovery, the benefit of HBOT or medical treatment may not have been accurately evaluated. There may be a bias effect caused by spontaneous recovery during HBOT as a salvage treatment or prolonged HBOT. However, in the case of HBOT as a salvage treatment, any benefit would not be significant because of the extended period between onset and therapy. For prolonged HBOT, spontaneous recovery during therapy cannot be distinguished from treatment benefits. Therefore, given a sufficient pooled sample size, it is reasonable to assume that the likelihood of such benefit would be evenly distributed in both groups.

Conclusions

For SSNHL, HBOT + MT was shown to be a more advantageous treatment option than MT alone. A benefit of HBOT was observed in patients with SSNHL who had severe to profound hearing loss at baseline and who underwent HBOT as a salvage treatment with a prolonged total HBOT duration. Further trials using well-defined indications and standardized protocols of HBOT are warranted.

ARTICLE INFORMATION

Accepted for Publication: July 12, 2018.

Published Online: September 27, 2018.
doi:10.1001/jamaoto.2018.2133

Author Contributions: Dr Rhee had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Drs Rhee and Hwang contributed equally to this work.

Concept and design: Rhee, Hwang, J.-S. Lee.
Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Rhee.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Rhee, Hwang, Park.

Supervision: J. M. Lee.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest, and none were reported.

REFERENCES

1. Stachler RJ, Chandrasekhar SS, Archer SM, et al; American Academy of Otolaryngology–Head and Neck Surgery. Clinical practice guideline: sudden hearing loss. *Otolaryngol Head Neck Surg*. 2012;146(3)(suppl):S1–S35. doi:10.1177/0194599812436449
2. Wie OB, Prupp AH, Tvete O. Unilateral deafness in adults: effects on communication and social interaction. *Ann Otol Rhinol Laryngol*. 2010;119(11):772–781.
3. Murphy-Lavoie H, Piper S, Moon RE, Legros T. Hyperbaric oxygen therapy for idiopathic sudden sensorineural hearing loss. *Undersea Hyperb Med*. 2012;39(3):777–792.

4. Gill AL, Bell CN. Hyperbaric oxygen: its uses, mechanisms of action and outcomes. *QJM*. 2004;97(7):385–395. doi:10.1093/qjmed/hch074
5. Appaia A, Pech A, Demard F. [The use of hyperbaric oxygen in oto-rhino-laryngology] [in French]. *Ann Otolaryngol Chir Cervicofac*. 1970;87(12):735–750.
6. Topuz E, Yigit O, Cinar U, Seven H. Should hyperbaric oxygen be added to treatment in idiopathic sudden sensorineural hearing loss? *Eur Arch Otorhinolaryngol*. 2004;261(7):393–396. doi:10.1007/s00405-003-0688-6
7. Cekin E, Cincik H, Ulubil SA, Gungor A. Effectiveness of hyperbaric oxygen therapy in management of sudden hearing loss. *J Laryngol Otol*. 2009;123(6):609–612. doi:10.1017/S0022215109004277
8. Cvorovic L, Jovanovic MB, Milutinovic Z, Arsovic N, Djeric D. Randomized prospective trial of hyperbaric oxygen therapy and intratympanic steroid injection as salvage treatment of sudden sensorineural hearing loss. *Otol Neurotol*. 2013;34(6):1021–1026. doi:10.1097/MAO.0b013e318297638a
9. Narozny W, Kuczkowski J, Kot J, Stankiewicz C, Sicko Z, Mikaszewski B. Prognostic factors in sudden sensorineural hearing loss: our experience and a review of the literature. *Ann Otol Rhinol Laryngol*. 2006;115(7):553–558. doi:10.1177/000348940611500710
10. Desloovere C, Knecht R, Germonpré P. Hyperbaric oxygen therapy after failure of conventional therapy for sudden deafness. *B-ENT*. 2006;2(2):69–73.
11. Satar B, Hidir Y, Yetiser S. Effectiveness of hyperbaric oxygen therapy in idiopathic sudden

hearing loss. *J Laryngol Otol*. 2006;120(8):665–669. doi:10.1017/S0022215106001769

12. Dunder K, Gumus T, Ay H, Yetiser S, Ertugrul E. Effectiveness of hyperbaric oxygen on sudden sensorineural hearing loss: prospective clinical research. *J Otolaryngol*. 2007;36(1):32–37. doi:10.2310/7070.2006.0061

13. Fujimura T, Suzuki H, Shiomori T, Udaka T, Mori T. Hyperbaric oxygen and steroid therapy for idiopathic sudden sensorineural hearing loss. *Eur Arch Otorhinolaryngol*. 2007;264(8):861–866. doi:10.1007/s00405-007-0272-6

14. Ohno K, Noguchi Y, Kawashima Y, Yagishita K, Kitamura K. Secondary hyperbaric oxygen therapy for idiopathic sudden sensorineural hearing loss in the subacute and chronic phases. *J Med Dent Sci*. 2010;57(2):127–132.

15. Alimoglu Y, Inci E, Edizer DT, Ozdilek A, Aslan M. Efficacy comparison of oral steroid, intratympanic steroid, hyperbaric oxygen and oral steroid + hyperbaric oxygen treatments in idiopathic sudden sensorineural hearing loss cases. *Eur Arch Otorhinolaryngol*. 2011;268(12):1735–1741. doi:10.1007/s00405-011-1563-5

16. Liu SC, Kang BH, Lee JC, et al. Comparison of therapeutic results in sudden sensorineural hearing loss with/without additional hyperbaric oxygen therapy: a retrospective review of 465 audiological controlled cases. *Clin Otolaryngol*. 2011;36(2):121–128. doi:10.1111/j.1749-4486.2011.02303.x

17. Yang CH, Wu RW, Hwang CF. Comparison of intratympanic steroid injection, hyperbaric oxygen and combination therapy in refractory sudden sensorineural hearing loss. *Otol Neurotol*. 2013;34(8):1411–1416. doi:10.1097/MAO.0b013e3182a1eb83

18. Capuano L, Cavaliere M, Parente G, et al. Hyperbaric oxygen for idiopathic sudden hearing loss: is the routine application helpful? *Acta Otolaryngol*. 2015;135(7):692-697. doi:10.3109/00016489.2015.1023355
19. Pezzoli M, Magnano M, Maffi L, et al. Hyperbaric oxygen therapy as salvage treatment for sudden sensorineural hearing loss: a prospective controlled study. *Eur Arch Otorhinolaryngol*. 2015; 272(7):1659-1666. doi:10.1007/s00405-014-2948-z
20. Psillas G, Ouzounidou S, Stefanidou S, et al. Hyperbaric oxygen as salvage treatment for idiopathic sudden sensorineural hearing loss. *B-ENT*. 2015;11(1):39-44.
21. Hosokawa S, Sugiyama KI, Takahashi G, et al. Hyperbaric oxygen therapy as adjuvant treatment for idiopathic sudden sensorineural hearing loss after failure of systemic steroids. *Audiol Neurootol*. 2017;22(1):9-14. doi:10.1159/000464096
22. Cavallazzi G, Pignataro L, Capaccio P. Italian experience in hyperbaric oxygen therapy for idiopathic sudden sensorineural hearing loss. In: Proceedings of the International Joint Meeting on Hyperbaric and Underwater Medicine. Bologna, Italy: Grafica Victoria; 1996:647-649.
23. Aslan I, Oysu C, Veyseller B, Baserer N. Does the addition of hyperbaric oxygen therapy to the conventional treatment modalities influence the outcome of sudden deafness? *Otolaryngol Head Neck Surg*. 2002;126(2):121-126. doi:10.1067/mhn.2002.121915
24. Ricciardiello F, Abate T, Pianese A, et al. Sudden sensorineural hearing loss: role of hyperbaric oxygen therapy. *Translational Med Rep*. 2017;1:6497.
25. Bennett MH, Kertesz T, Perleth M, Yeung P, Lehm JP. Hyperbaric oxygen for idiopathic sudden sensorineural hearing loss and tinnitus. *Cochrane Database Syst Rev*. 2012;10:CD004739.
26. Saesen K, Loos E, Montagna C, Vanbrabant T, Goedhuys R, Lemkens N. Hyperbaric oxygen therapy in idiopathic sudden sensorineural hearing loss. *B-ENT*. 2017;13(2):105-112. <https://www.b-ent.be/articles/hyperbaric-oxygen-therapy-in-idiopathic-sudden-sensorineural-hearing-loss>. Accessed August 14, 2018.
27. Normand SL. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med*. 1999;18(3):321-359. doi:10.1002/(SICI)1097-0258(19990215)18:3<321::AID-SIM28>3.0.CO;2-P
28. Ahmed I, Sutton AJ, Riley RD. Assessment of publication bias, selection bias, and unavailable data in meta-analyses using individual participant data: a database survey. *BMJ*. 2012;344:d7762. doi:10.1136/bmj.d7762
29. Wei BP, Stathopoulos D, O'Leary S. Steroids for idiopathic sudden sensorineural hearing loss. *Cochrane Database Syst Rev*. 2013;(7):CD003998.
30. Nagahara K, Fisch U, Yagi N. Perilymph oxygenation in sudden and progressive sensorineural hearing loss. *Acta Otolaryngol*. 1983; 96(1-2):57-68. doi:10.3109/00016488309132875
31. Kim SA, Ahn JH. Clinical application of hyperbaric oxygen in treatment of idiopathic sudden sensorineural hearing loss. *Korean J Otorhinolaryngol Head Neck Surg*. 2016;59(7): 490-494. doi:10.3342/kjorl-hns.2016.59.7.490
32. Lawrence R, Thevasagayam R. Controversies in the management of sudden sensorineural hearing loss: an evidence-based review. *Clin Otolaryngol*. 2015;40(3):176-182. doi:10.1111/coa.12363
33. Plafki C, Peters P, Almeling M, Welslau W, Busch R. Complications and side effects of hyperbaric oxygen therapy. *Aviat Space Environ Med*. 2000;71(2):119-124.